Dynamic Perturbation Spreading in Networks

Malte Schröder^(D), Xiaozhu Zhang, Justine Wolter, and Marc Timme

Abstract-Understanding how local perturbations induce the transient dynamics of a network of coupled units is essential to control and operate such systems. Often a perturbation initiated in one unit spreads to other units whose dynamical state they transiently alter. The maximum state changes at those units and the timings of these changes constitute key characteristics of such transient response dynamics. However, even for linear dynamical systems it is not possible to analytically determine time and amplitude of the maximal response of a unit to a perturbation. Here, we propose to extract approximate peak times and amplitudes from effective expectation values used to characterize the typical time and magnitude of the response of a unit by interpreting the system's response as a probability distribution over time. We derive analytic estimators for the peak response based on these expectation value measures in linearized systems operating close to a stable fixed point. These estimators can be expressed in terms of the inverse of the system's Jacobian. We obtain identical results with different approximations for the response dynamics, indicating that these estimators become exact in the limit of weak coupling. Furthermore, the results suggest that perturbations spread ballistically in networks with diffusive coupling.

Index Terms—Network dynamics, perturbations, perturbation spreading.

I. INTRODUCTION

TRANSIENT collective dynamics plays an important role in a wide range of systems from social and biological systems where ideas or diseases spread [1]–[6] to the stability of large scale infrastructure and supply networks such as power grids [7]–[10]. These systems generically operate near a fixed point and are naturally subject to perturbations, for example an outbreak of an infection or fluctuations of the power consumption and production [11]. In their simplest setting, such perturbations initially affect only a single unit and spread through the network, transiently affecting other units at different times and with different strengths [12], [13].

Despite the importance of these spreading and propagation processes, no general answer exists for when or how strongly a

The authors are with the Chair for Network Dynamics of the Center for Advancing Electronics Dresden (cfaed), the Institute for Theoretical Physics and the Cluster of Excellence Physics of Life, TU Dresden, 01062 Dresden, Germany (e-mail: malte.schroeder@tu-dresden.de; xiaozhu.zhang@tu-dresden.de; justine. wolter@stud.uni-goettingen.de; marc.timme@tu-dresden.de).

This paper has supplementary downloadable material available at http://ieeexplore.ieee.org, provided by the authors.

Digital Object Identifier 10.1109/TNSE.2019.2901582

unit is affected by an initial perturbation. Traditional measures to characterize these transient responses are the time and magnitude of the maximal (peak) response. However, even in linearized systems, computing such measures typically involves the solution of transcendental equations making exact analytical predictions impossible.

A recently introduced idea [14] is interpreting the deterministic transient responses as probability densities in time. The resulting "effective expectation values" constitute characteristic response measures (different from traditional ones) that are computable analytically in linearized systems in terms of the inverse effective coupling matrix.

In this article we derive analytic estimators for the peak response time and amplitude based on these expectation value response measures in linearized systems affected by perturbations around a stable fixed point. We approximate the response dynamics with multiple different functions that qualitatively reproduce it. For each approximation function, we analytically derive both, the response strength and timing as calculated from the expectation values and the amplitude and timing of maximal response. Comparing these results we find analytic estimators for the actual peak response time and magnitude in terms of the inverse Jacobian of the linearized system. We illustrate that such estimators become exact in the limit of weak coupling, independent of the topology of the coupling network.

II. PERTURBATIONS IN NETWORK DYNAMICAL SYSTEMS

Consider a general network dynamical system

$$\frac{\mathrm{d}\mathbf{y}}{\mathrm{d}t} = \mathbf{F}(\mathbf{y}) \tag{1}$$

consisting of N coupled units i with internal state $y_i(t) = [\mathbf{y}(t)]_i$ operating close to a stable fixed point $\mathbf{y}^* \in \mathbb{R}^N$. Small perturbations to this state and their impact across the network (Fig. 1) are described by the linearized dynamics $d\mathbf{x}/dt = M\mathbf{x}$ where $\mathbf{x}(t) = \mathbf{y}(t) - \mathbf{y}^*$ and $M = d\mathbf{F}/d\mathbf{y}|_{\mathbf{y}=\mathbf{y}^*}$ is the Jacobian. The diagonal elements of M describe the internal dynamics of the individual units while the off-diagonal elements describe the coupling between the units.

In general, the impact of a perturbation on a unit and how a perturbation spreads through the network can be measured in different ways: for models of epidemic spreading, describing an outbreak across different populations coupled by a transportation network, arrival times are often defined by measuring the first time $t_{\text{arrival}} = \min\{t : x_i(t) \ge \epsilon\}$ when the number of infected individuals exceeds a given threshold ϵ . For stochastic epidemic spreading the connection to random walk processes allows predictions of the arrival times of the perturbation [1],

2327-4697 © 2019 IEEE. Personal use is permitted, but republication/redistribution requires IEEE permission. See https://www.ieee.org/publications/rights/index.html for more information.

Manuscript received December 4, 2018; revised January 28, 2019; accepted February 21, 2019. Date of publication February 26, 2019; date of current version September 2, 2020. This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2068 – 390729961 – Cluster of Excellence Physics of Life and the Cluster of Excellence Center for Advancing Electronics at TU Dresden and the German Federal Ministry for Research and Education (BMBF grants no. 03SF0472F and 03EK3055F). Recommended for acceptance by J. G. Barajas Ramírez. (*Corresponding author: Malte Schröder.*)

¹⁰¹⁹



Fig. 1. **Typical response to a localized perturbation.** (a) A small network dynamical system with N = 6 units and E = 9 directed interactions. (b) Responses $x_i(t)$ of all units *i* after an initial perturbation $x_1(0) = 1$ at unit k = 1 (dark yellow) with homogeneous coupling strength $\alpha = 1$ and internal dynamics $\beta = 1$. The perturbation spreads through the network and transiently affects all other units before the system returns to the stable operating state $x_i^* = 0$ for $t \to \infty$. (c, d) For small times $t \to 0$ the activity $x_i(t)$ of unit *i* grows polynomially as t^d with an exponent d = i - 1, the graph-theoretical distance from the initially perturbed unit. For large times $t \to \infty$ the activity of all units decays exponentially with exponent $-\beta$.

[3], [4], [15], [16]. For other spreading processes the total impact of the perturbation at a given unit or the maximal deviation from the operating point is of interest [12], [13], [17]–[20]. Interestingly, even for the simple deterministic linear system $d\mathbf{x}/dt = M\mathbf{x}$ described above, these measures cannot be easily evaluated analytically. The underlying reason is that calculating peak positions or threshold crossing times typically involves solving transcendental equations of the form $a_1 \exp(\lambda_1 t) + a_2 \exp(\lambda_2 t) + \ldots = c$ for the time *t*, where λ_j are the eigenvalues of *M*.

Recently, a complementary approach was introduced to characterize the impact of a perturbation in such linearized systems in terms of expectation values of effective probability distributions [14]. This approach works as follows: We first normalize each response trajectory and interpret the result as a probability density over time. We then quantify the arrival time and impact of a perturbation by expectation values and higher order moments with respect to this effective probability distribution.

In the following we assume that the internal dynamics of each unit is described by exponential decay with a rate $\beta_i > 0$ $(\mathbf{x} = \mathbf{0} \text{ is a stable fixed point)}$ and the coupling between the units is diffusive with coupling strength $\alpha_{ij} \ge 0$, such that $M_{ij} = \alpha_{ij}$ (for $i \ne j$), $\alpha_{ij} = 0$ if unit *i* is not directly affected by unit *j* and $M_{ii} = -\beta_i - \sum_j \alpha_{ij}$. In this case the response of each unit $x_i(t)$ to an initial perturbation $x_i(0) = (\mathbf{x}_0)_i = \delta_{ik}$ at a single unit *k* is guaranteed to be positive, $x_i(t) > 0$ for all times t > 0 (see Fig. 1). Appropriate normalization of the response trajectories $x_i(t) = [\exp(Mt)\mathbf{x}_0]_i$ by the total response

$$Z_i = \int_0^\infty x_i(t) \mathrm{d}t = -\left(M^{-1} \mathbf{x_0}\right)_i \tag{2}$$

then allows the interpretation of the trajectories as if they were probability densities over time, $\rho_i(t) = x_i(t)/Z_i$. From this perspective, expectation values of time with respect to the probability distribution characterize the impact of the perturbation at different units (see Fig. 2) with simple analytic expressions [14]. For example, the expectation value

$$\langle t \rangle_i = \int_0^\infty t \rho_i(t) \mathrm{d}t = -\frac{(M^{-2} \mathbf{x}_0)_i}{(M^{-1} \mathbf{x}_0)_i} \tag{3}$$

describes the characteristic response time (not the peak time) when the perturbation impacts unit *i*. Similarly, the typical duration of the perturbation is measured in terms of the standard deviation σ_i and its magnitude by the quotient H_i of total response Z_i and the standard deviation

$$\sigma_i = \sqrt{\int_0^\infty (t - \langle t \rangle_i)^2 \rho_i(t) dt}$$

$$= \sqrt{\frac{2(M^{-3} \mathbf{x}_0)_i}{(M^{-1} \mathbf{x}_0)_i} - \left(\frac{(M^{-2} \mathbf{x}_0)_i}{(M^{-1} \mathbf{x}_0)_i}\right)^2}$$
(4)

$$H_{i} = \frac{Z_{i}}{\sigma_{i}} = \frac{\left((M^{-1}\mathbf{x}_{0})_{i} \right)^{2}}{\sqrt{2(M^{-3}\mathbf{x}_{0})_{i}(M^{-1}\mathbf{x}_{0})_{i} - \left((M^{-2}\mathbf{x}_{0})_{i} \right)^{2}}} .$$
 (5)

These quantities are illustrated in Fig. 2 together with the numerically determined peak response values t_i^{peak} and $x_i^{\text{peak}} = x_i(t_i^{\text{peak}})$ for the example system from Fig. 1. As also demonstrated previously [14], the characteristic response times and response magnitudes [Eq. (3) and (5)] appear to show the same scaling as the actual peak time t_i^{peak} and the maximal response $x_i(t_i^{\text{peak}})$, that means they accurately describe the *relative* impact of the perturbations at different units. However, if interpreted as estimators for the peak response values they are clearly biased and do not provide a good quantitative description of the *absolute* impact.

In general, for unimodal distributions as we observe for the typical response trajectories, some conditions on the relationship between mean (expectation value) and mode (position of the maximum) are known. For example, if the distribution has positive skewness (as the response trajectories), we typically have $\langle t \rangle_i \geq t_i^{\text{peak}}$. Unimodal distributions also satisfy the



Fig. 2. Expectation values quantify the relative impact of the perturbation. (a) Interpretation of the response trajectories as probability densities over time yields the characteristic response measures introduced in [14] based on the effective expectation values defined in Eq. (2 - 5). The example shows the response $x_6(t)$ of unit 6 (dark green, compare Fig. 1) and the corresponding characteristic response measures (red). (b,c) These measures accurately characterize the *relative* impact of the perturbation and show the same scaling as the actual peak response times and magnitudes, illustrated here for the small example shown in Fig. 1 where d = i - 1 is the graph-theoretical distance from the perturbed unit. The response times seem to be biased additively, the response strengths multiplicatively. This observed systematic difference suggests that an adjustment is possible to obtain estimators for the *absolute* peak response values based on the characteristic (expectation value) response measures.

condition $\left|t_i^{\text{peak}} - \langle t \rangle_i\right| \leq \sqrt{3}\sigma_i$ [21]. Unfortunately, no exact connection between mean and mode for general distributions, and thereby for general $x_i(t)$, exists.

For the specific class of (initially algebraically increasing and then exponentially decaying) responses $x_i(t)$ in linearized systems, we here establish two connections between the characteristic response values and the actual peak values. Specifically, the examples in [14] already suggest that the relation between the actual peak measures and the characteristic response measures is systematic and largely independent of the structure of the interaction network, as also illustrated in Fig. 2 (b,c). As illustrated in Fig. 2(b), we observe an approximately constant shift between $\langle t \rangle_i$ and t_i^{peak} . This suggests an additive adjustment c_T to estimate the actual peak time

$$\operatorname{Est}\left[t^{\operatorname{peak}}\right]_{i} = \langle t \rangle_{i} + c_{T} \,. \tag{6}$$

Similarly, as illustrated in Fig. 2(c) we also observe a constant multiplicative factor between H_i and x_i^{peak} (note the logarithmic axis), suggesting a multiplicative adjustment c_H such that

$$\operatorname{Est}\left[x^{\operatorname{peak}}\right]_{i} = c_{H} H_{i} \,. \tag{7}$$

In the following we analytically derive these adjustments from approximate response functions and show that they results in the same form of adjustment. We use these calculations to determine the constants c_T and c_H and define the estimators for the actual peak response based on approximating model trajectories. For different classes of model functions that recover the qualitative shape and the asymptotic behavior of the response dynamics we calculate exact characteristic and peak response values as an explicit function of the interaction network described by M. We use these expressions to convert the characteristic response measures resulting from effective expectation values into estimators for the peak values for these approximating functions and thereby for the real response dynamics.

III. ESTIMATORS FOR THE PEAK RESPONSE

The asymptotic behavior of the units' responses is given by polynomial growth for $t \to 0$ and by exponential decay for $t \to \infty$ [compare Fig. 1(c,d)]. Motivated by these known asymptotic scaling regimes, we illustrate the approach to calculate the constants c_T and c_H using a simple family of approximating functions

$$\tilde{x}(t) = \tilde{A} t^d \exp(-\tilde{\beta}t) \tag{8}$$

that roughly capture the qualitative behavior of the response dynamics. Since the constant factor \tilde{A} does not change the peak position or the factors c_T and c_H , we set $\tilde{A} = 1$ in the following. We now first determine the remaining parameters \tilde{d} and $\tilde{\beta}$ of this approximate response function.

To understand the asymptotic behavior of the units for small times, $t \rightarrow 0$, we consider the formal solution

$$\mathbf{x}(t) = \exp(Mt)\mathbf{x}_0$$

= $\mathbf{x}_0 + tM\mathbf{x}_0 + 1/2 t^2 M^2 \mathbf{x}_0 + \dots$ (9)

For a perturbation at a single unit k with $x_k(0) = 1$ this expression reduces to the matrix elements $x_i(t) = \delta_{ik} + tM_{ik} + 1/2 t^2 (M^2)_{ik} + \ldots$. The matrix M is directly related to the adjacency matrix of the interaction network (with additional entries along the diagonal). Thus, for networks with homogenous coupling strengths $\alpha_{ij} = \alpha$, the element $(M^n)_{ik}$ is proportional to the number of paths from k to i of length n [22]. If we denote the (shortest path) distance from the initially perturbed unit k to unit i as $d \in \mathbb{N}$, all elements $(M^n)_{ik} = 0$ for n < d since there are no paths of length n < d. This means that the first non-zero term in the response of unit i is given by $x_i(t) = 1/d!(M^d)_{ik}t^d + \mathcal{O}(t^{d+1})$ as $t \to 0$. The same argument holds for networks with heterogenous coupling strengths α_{ij} . The entry $(M^n)_{ik}$ is then given by the sum over all weighted paths of length n.

For large times, $t \to \infty$, we consider the eigenvalues $-\lambda_i$ of M. We explicitly write them as $-\lambda_i$ to signify that all

eigenvalues have negative real part $\operatorname{Re}[-\lambda_i] < 0$ since M describes the dynamics around an asymptotically stable fixed point. We label the eigenvalues such that $\operatorname{Re}[-\lambda_N] \leq ... \leq \operatorname{Re}[-\lambda_2] \leq \operatorname{Re}[-\lambda_1]$. The response in terms of these eigenvalues is then given by

$$x_{i}(t) = [\mathbf{x}(t)]_{i} = [\exp(Mt)\mathbf{x}_{0}]_{i}$$

= $c_{1} \exp(-\lambda_{1}t) + c_{2} \exp(-\lambda_{2}t) +$
= $\exp(-\lambda_{1}t)[c_{1} + c_{2} \exp((\lambda_{1} - \lambda_{2})t) +],$
(10)

with constants c_i depending on the initial conditions \mathbf{x}_0 . For undirected networks (symmetric M) $c_j = \left(\mathbf{v}_j^T \mathbf{x}_0\right) \left[\mathbf{v}_j\right]_i$ where $\mathbf{v}_1, \mathbf{v}_2$, denote the orthogonal eigenvectors of M corresponding to the (real) eigenvalues $-\lambda_1, -\lambda_2$,. For large $t \to \infty$ the first term dominates. Thus, the response at all units is given by $x_i(t) = c_1 \exp(-\lambda_1 t) + \mathcal{O}[\exp(-\lambda_2 t)]$ with magnitude $\mathcal{O}[\exp(-\lambda_1]t)]$. For notational convenience, we do the following calculations for undirected networks with real eigenvalues and drop the real part notation, writing only $-\lambda_1$.

Matching the above considerations for large and small t, defining the parameters of $\tilde{x}(t)$ as $\tilde{d} = d$ and $\tilde{\beta} = \lambda_1$, Eq. (8) becomes

$$\tilde{x}(t) = t^d \exp(-\lambda_1 t) \,. \tag{11}$$

For this approximation for the response of a unit at (shortest path) distance d to the initial perturbation we now calculate both the typical response measures [Eq. (3) and (5)] as well as the true peak response values analytically. The normalization factor \tilde{Z} [Eq. (2)] is

$$\tilde{Z} = \int_0^\infty \tilde{x}(t) \mathrm{d}t = \frac{d!}{\lambda_1^{d+1}} , \qquad (12)$$

and we define $\tilde{\rho}(t) = \tilde{x}(t)/\tilde{Z}$. Here and in the following we drop the indices of the response function denoting the dependence on the unit *i* (and the initially perturbed unit *k*). These dependencies become explicit by noting that the graph-theoretical distance *d* is a function of *i* and *k*. The higher order moments follow analogously to Eq. (12) by definition [see Eq. (3) and (5)] as

$$\langle \tilde{t} \rangle = \frac{d+1}{\lambda_1} \tag{13}$$

$$\tilde{\sigma} = \sqrt{\langle \tilde{t}^2 \rangle - \langle \tilde{t} \rangle^2} = \sqrt{\frac{d+1}{\lambda_1^2}} \tag{14}$$

$$\tilde{H} = \frac{\tilde{Z}}{\tilde{\sigma}} = \frac{d!}{\lambda_1^d \sqrt{d+1}}$$

$$\sim \sqrt{2\pi} \left(\frac{d}{\lambda_1}\right)^d \exp(-d) \quad \text{as} \quad d \to \infty \,,$$
(15)

where the last line denotes the asymptotic behavior for large distances $d \rightarrow \infty$ (see Supplemental Material for detailed derivations).

To obtain the peak response measures we determine the maximum $\frac{d}{dt}\tilde{x}(t)|_{t=\tilde{t}^{peak}}=0$ and find

$$\tilde{t}^{\text{peak}} = \frac{d}{\lambda_1} \tag{16}$$

and consequently

$$\widetilde{x}^{\text{peak}} = \widetilde{x}(\widetilde{t}^{\text{peak}}) = \left(\frac{d}{\lambda_1}\right)^d \exp(-d).$$
(17)

Comparing Eq. (16) to (13) and Eq. (17) to (15) suggests the bias corrections

$$c_T = \tilde{t}^{\text{peak}} - \langle \tilde{t} \rangle = -\frac{1}{\lambda_1} \tag{18}$$

and

$$c_{H} = \frac{\tilde{x}^{\text{peak}}}{\tilde{H}} = \frac{\sqrt{d+1} d^{d}}{\exp(d) d!}$$

$$= \frac{1}{\sqrt{2\pi}} + \mathcal{O}(d^{-1}),$$
(19)

where the last line describes the asymptotic behavior for large distances d. Note that these factors are independent of the origin of the perturbation k or the specific unit i for large distances but only depend on the network structure through the largest eigenvalue $-\lambda_1$. Analogously, we now convert the characteristic response measures for the original trajectories $x_i(t)$ to the estimators for the peak height and position

$$\operatorname{Est}\left[t^{\operatorname{peak}}\right]_{i} = \langle t \rangle_{i} - \frac{1}{\lambda_{1}} = -\frac{(M^{-2}\mathbf{x}_{0})_{i}}{(M^{-1}\mathbf{x}_{0})_{i}} - \frac{1}{\lambda_{1}} \qquad (20)$$

$$\operatorname{Est}\left[x^{\operatorname{peak}}\right]_{i} = \frac{\sqrt{d+1} d^{d}}{\exp(d) d!} H_{i}$$

$$= \frac{H_{i}}{\sqrt{2\pi}} + \mathcal{O}(d^{-1})$$
(21)

where d = d(i, k) is the graph theoretical distance from the perturbed unit k to unit i. Consequently, this also suggests the new definition of the typical response duration as $\Delta t_i = Z_i/\text{Est}$ $[x(t^{\text{peak}})]_i = \sqrt{2\pi} \sigma_i + \mathcal{O}(d^{-1})$, illustrated in Fig. 3.

IV. UNIVERSAL BALLISTIC SPREADING FOR WEAK COUPLING?

The calculations above do not mathematically imply that these results should extend to the real response dynamics. Specifically, the limiting behavior of $\tilde{x}(t)$ is not purely exponential for $t \to \infty$ but scales as $t^d \exp(-\lambda_1 t)$. Beyond numerical validation of the results, we compute the same adjustments for other approximating functions that *do* exhibit the correct asymptotic scaling for both $t \to 0$ and $t \to \infty$. Importantly, comparing the results of all of these calculations, we find *identical* adjustments as above in the limit of weak coupling $\alpha/\lambda_1 \to 0$ (see Supplemental Material for details and calculations). Numerical analysis, illustrated in Fig. 4, supports that estimator errors indeed decay to zero for weak coupling $\alpha/\lambda_1 \to 0$. Together, these results suggest that the adjustments we derived above are universal in this limit.



Fig. 3. Adjusted expectation values. (a) Illustration of the adjusted expectation value quantifiers (compare Fig 2). (b,c) With the adjustments derived in the main text the expectation values [Eq. (2 - 5)] are converted into more accurate estimators [Eq. (20) and (21)] for the actual time and especially the peak magnitude of the response.

Moreover, both the peak response time as well as the characteristic response times increase linearly with the distance d in this limit for all families of approximating response functions. This indicates that the spreading of the perturbation is ballistic, even though the coupling is diffusive. This observation is in line with heuristic predictions for different dynamics such as diseases spreading in transportation networks [1]–[4], [6]. In these models the mobility rate (coupling strength) is typically much slower then



Fig. 4. Exact estimators in the limit of weak coupling. (a) Relative error of the estimated peak height Eq. (21). (b) Absolute error of the estimated peak time Eq. (20). Both errors disappear for large λ_1/α , that means for weak coupling $\alpha/\lambda_1 \rightarrow 0$. Each point indicates one observation in R = 100 connected Poisson random networks with N = 20 units and M = 40 links with fixed $\lambda_1 = \beta = 1$ and variable coupling strength α , regardless of the distance of the unit to the origin of the perturbation.

the internal dynamics of the individual units (weak coupling limit) and the observed arrival time increases linearly with the (effective) distance of a unit to the original outbreak location.

V. ACCURATE ESTIMATION ACROSS NETWORK TOPOLOGIES

We numerically test the accuracy of the estimators across different network topologies for fixed α/λ_1 . We perturb each unit in the network once and record the resulting typical response times and magnitudes as well as peak values. For simplicity we use constant coupling strengths $\alpha_{ij} \in \{0, \alpha\}$ and identical internal dynamics $\beta_i = \beta = \lambda_1 = 1$ in these examples. However, this is not a necessary condition for our results to hold as the derivation given above holds for general matrices M, assuming only $x_i(t) > 0$ for all t > 0 and $x_i(t) \to 0$ as $t \to \infty$.

Figure 5 shows the results for Poisson random networks (narrow degree distribution, small diameter), Barabasi-Albert scale free random networks (broad degree distribution, small diameter) and random geometrically embedded networks (narrow degree distribution, large diameter) for $\alpha/\lambda_1 = 1$. The adjustment systematically improves the estimate compared to the characteristic response values but is still not exact, as expected for non-zero α/λ_1 . Specifically, the peak time is typically underestimated. The estimate of the peak height becomes more accurate for large distances (small x^{peak}). Figure 6 shows the same simulations with weaker coupling $\alpha/\lambda_1 = 0.1$. As expected from the analytical calculations, the estimates agree much better with the exact peak values. Additional results for absolute and relative errors of the estimators are shown in the Supplemental Material.

We specifically note, that all assumptions in the derivations presented above are satisfied also for directed networks or networks with heterogeneous coupling strengths. As seen in Fig. 6 (b), heterogeneous network structures (and similarly heterogeneous coupling strengths) cause larger fluctuations in the estimations. The reason is the existence of multiple short paths or stronger coupling along these paths in such networks. However, the analytic results remain correct. Most importantly, the estimators become exact in the limit of weak coupling, independent of the network topology or coupling strength distribution.



Fig. 5. Improved estimation of peak values across network topologies. The top row shows examples of the network topologies of (a) Poisson random networks, (b) Barabasi-Albert scale free random networks and (c) geometrically embedded random networks with periodic boundaries. The middle row shows the resulting $\text{Est}[x(t^{\text{peak}})]$ [blue, Eq.(21)] versus the true peak height x^{peak} , the bottom row shows the corresponding results for $\text{Est}[t^{\text{peak}}]$ [blue, Eq.(20)]. Points on the diagonal indicate perfect agreement of the estimated peak time or height with the actual peak time or height. Both estimators improve the prediction of the actual peak values compared to the raw expectation values (red). In both cases the estimators become more accurate for larger distances (smaller x^{peak} and larger t^{peak}). All networks consist of N = 200 units with E = 400 undirected interactions. The simulation parameters are $\beta = \lambda_1 = 1$ and $\alpha/\lambda_1 = 1$ in all three cases. The plots shows results for R = 10 different realizations of the network structure where every unit was perturbed once, for a total of 400000 measurements.

VI. CONCLUSION

Understanding the propagation of perturbation-induced signals in networked systems helps to predict, control and mitigate their impact in a range of processes in biology and engineering, from epidemic spreading of diseases to the impact of load shedding or infrastructure outages in electric power grids. Among the fundamental questions are when and how strongly perturbations initiated at some unit in a network reach other units. So far, it has been impossible to analytically estimate timing and strengths of such signals as an explicit function of the underlying base state of the system and the network's interaction topology. These limitations hold even for linear deterministic systems because the equations determining peak timing and strength are transcendental and as such mathematically intractable.

A recent proposal [14] suggests to take a complementary perspective and predicts characteristic arrival times and strengths not in terms of peak times and amplitudes but in terms of expectation values that result from interpreting the deterministic trajectory of a unit's response as a probability density. This approach yields characteristic arrival times and strengths as



Fig. 6. Accurate estimation of peak values across network topologies with weak coupling. The top row shows examples of the network topologies of (a) Poisson random networks, (b) Barabasi-Albert scale free random networks and (c) geometrically embedded random networks with periodic boundaries. The middle row shows the resulting $\text{Est}[x(t^{\text{peak}})]$ [blue, Eq.(21)] versus the true peak height x^{peak} , the bottom row shows the corresponding results for $\text{Est}[t^{\text{peak}}]$ [blue, Eq.(21)] versus the true peak height x^{peak} , the bottom row shows the corresponding results for $\text{Est}[t^{\text{peak}}]$ [blue, Eq.(20)]. Points on the diagonal indicate perfect agreement of the estimated peak time or height with the actual peak time or height. Both estimators more accurately predict the actual peak values compared to the raw expectation values (red). In both cases the estimators become more accurate for larger distances (smaller x^{peak} and larger t^{peak}). All networks consist of N = 200 units with E = 400 undirected interactions. The simulation parameters are $\beta = \lambda_1 = 1$ and $\alpha/\lambda_1 = 0.1$ in all three cases. The estimators are more accurate for this weaker coupling (compared to those in Fig. 5 for $\alpha = 1$). The plots show results for R = 10 different realizations of the network structure where every unit was perturbed once, for a total of 400000 measurements.

explicit functions of the inverse of the Jacobian matrix that in turn encodes both the base operating state and the interaction topology. However, these characteristic quantities are not intended to predict peak times and amplitudes – and if interpreted as such, exhibit large errors. So it still remains unclear how to explicitly quantify peak times and amplitudes.

Here we connect the two sets of quantifiers and derive approximate analytical estimators for the absolute peak positions and heights of the responses in terms of quantifiers based on expectation values. We employ qualitative approximations of the response functions mimicking the asymptotic behavior both for small and large times. The resulting estimators enable approximate predictions of the peak timings and heights across network topologies. Interestingly, in the weak coupling regime (see Supplemental Material), i.e. asymptotically as $\alpha/\lambda_1 \rightarrow 0$, the predictions become identical across all specific approximating functions tested, suggesting universality. Outside the asymptotic regime, i.e. for stronger coupling, the adjusted estimators seem to systematically underestimate the peak response values. Together with the expressions for the characteristic response measures derived by Wolter et al. in terms of expectation values [14], these results provide an analytic framework for predicting the impact of perturbations on any unit in any network operating close to a stable fixed point in the limit of weak coupling. Our results on deterministic systems are thereby complementing the analyses for specific models of disease spreading [1]–[4], [6]. They moreover suggest that in the asymptotic regime of weak coupling, perturbations spread ballistically through the network, even though the coupling is diffusive. Further work must show how details of the local network topology affect the accuracy of the predictions and how the results can be extended to allow also accurate predictions for stronger coupling and at close distances.

ACKNOWLEDGEMENTS

The authors would like to thank R. Schmidt and B. Friedrich for helpful discussions.

REFERENCES

- A. Gautreau, A. Barrat, and M. Barthélemy, "Arrival time statistics in global disease spread," *J. Statist. Mech.*, vol. 2007, no. 09, 2007, Art. no. L09001.
- [2] A. Gautreau, A. Barrat, and M. Barthelemy, "Global disease spread: Statistics and estimation of arrival times," *J. Theor. Biol.*, vol. 251, no. 3, pp. 509–522, 2008.
- [3] D. Brockmann and D. Helbing, "The hidden geometry of complex, network-driven contagion phenomena," *Science*, vol. 342, no. 6164, pp. 1337–1342, 2013.
- [4] F. Iannelli, A. Koher, D. Brockmann, P. Hövel, and I. M. Sokolov, "Effective distances for epidemics spreading on complex networks," *Phys. Rev. E*, vol. 95, Jan. 2017, Art. no. 012313.
- [5] C. Kirst, M. Timme, and D. Battaglia, "Dynamic information routing in complex networks," *Nat. Commun.*, vol. 7, 2016, Art. no. 11061.
- [6] L. M. Chen, M. Holzer, and A. Shapiro, "Estimating epidemic arrival times using linear spreading theory," *Chaos*, vol. 28, no. 1, 2018, Art. no. 013105.
- [7] D. Witthaut, M. Rohden, X. Zhang, S. Hallerberg, and M. Timme, "Critical links and nonlocal rerouting in complex supply networks," *Phys. Rev. Lett.*, vol. 116, 2016, Art. no. 138701.
- [8] D. Manik et al., "Network susceptibilities: Theory and applications," Phys. Rev. E, vol. 95, Jan. 2017, Art. no. 012319.
- [9] S. Kettemann, "Delocalization of disturbances and the stability of AC electricity grids," *Phys. Rev. E*, vol. 94, Dec. 2016, Art. no. 062311.
- [10] P. J. Menck, J. Heitzig, J. Kurths, and H. J. Schellnhuber, "How dead ends undermine power grid stability," *Nat. Commun.*, vol. 5, 2014, Art. no. 3969.
- [11] B. Schäfer, C. Beck, K. Aihara, D. Witthaut, and M. Timme, "Non-Gaussian power grid frequency fluctuations characterized by Lévy-stable laws and superstatistics," *Nat. Energy*, vol. 3, p. 119–126, 2018.
- [12] C. Hens, U. Harush, R. Cohen, and B. Barzel, "Predicting the patterns of spatio-temporal signal propagation in complex networks," 2018, arXiv:1801.08854.
- [13] M. Timme and J. Nagler, "Network science propagation patterns unravelled," *Nat. Phys.*, vol. 15, pp. 308–309, 2019.
- [14] J. Wolter, B. Lünsmann, X. Zhang, M. Schröder, and M. Timme, "Quantifying transient spreading dynamics on networks," *Chaos*, vol. 28, no. 6, 2018, Art. no. 063122.
- [15] L. A. Braunstein, S. V. Buldyrev, R. Cohen, S. Havlin, and H. E. Stanley, "Optimal paths in disordered complex networks," *Phys. Rev. Lett.*, vol. 91, Oct 2003, Art. no. 168701.
- [16] M. Roosta, "Routing through a network with maximum reliability," J. Math. Anal. Appl., vol. 88, no. 2, pp. 341–347, 1982.

- [17] T. Kittel, J. Heitzig, K. Webster, and J. Kurths, "Timing of transients: Quan tifying reaching times and transient behavior in complex systems," *New J. Phys.*, vol. 19, no. 8, 2017, Art. no. 083005.
- [18] B. K. Poolla, S. Bolognani, and F. Dörfler, "Optimal placement of virtual inertia in power grids," *IEEE Trans. Autom. Control*, vol. 62, no. 12, pp. 6209–6220, Dec. 2017.
- [19] M. Tyloo, T. Coletta, and P. Jacquod, "Robustness of synchrony in complex networks and generalized Kirchhoff indices," *Phys. Rev. Lett.*, vol. 120, Feb 2018, Art. no. 084101.
- [20] F. Hellmann, P. Schultz, C. Grabow, J. Heitzig, and J. Kurths, "Survivability of deterministic dynamical systems," *Sci. Rep.*, vol. 6, 2016, Art. no. 29654.
- [21] N. L. Johnson and C. A. Rogers, "The moment problem for unimodal distributions," Ann. Math. Statist., vol. 22, pp. 433–439, 1951.
- [22] X. Zhang, "Dynamic responses of networks under perturbations: Solutions, patterns and predictions," Ph.D. dissertation, Georg August Universität Göttingen, Göttingen, Germany, 2018.

Malte Schröder received the B.Sc., M.Sc. and Ph.D. degrees in physics from the Georg August University, Göttingen, Germany, and the IMPRS for Physics of Biological and Complex Systems, Max Planck Institute for Dynamics and Self-Organization, Göttingen, in 2012, 2014, and 2018, respectively. He is a Postdoctoral Researcher with the Chair for Network Dynamics, Institute for Theoretical Physics and the Center for Advancing Electronics Dresden, the Technical University Dresden. His research interests include spreading and synchronization in network dynamical systems, statistical physics of network formation and restructuring processes and the physics of transportation on networks. He is a member of German Physical Society.

Xiaozhu Zhang received the Ph.D. degree in physics from the Georg August University, Göttingen, Germany, and the IMPRS for Physics of Biological and Complex Systems, the Max Planck Institute for Dynamics and Self-Organization, Göttingen. She is a Postdoctoral Researcher with the Chair for Network Dynamics, Institute for Theoretical Physics and the Center for Advancing Electronics Dresden, Technical University Dresden. Her research interests include perturbation spreading and pattern formation on networks of dynamical systems, and collective dynamics of power grids and transport networks. She is a Guest Editor of the Focus Issue: Dynamics of Modern Power Grids of "Chaos: An Interdisciplinary Journal of Nonlinear Science" and has been one of the organizers of the mini-symposium "Structure and dynamics of future energy systems: power grids as complex dynamical systems" at Dynamics Days Europe 2018. She is a member of German Physical Society and Complex Systems Society.

Justine Wolter received the B.Sc. degree in physics in 2017 with a thesis on the prediction of perturbation spreading across networks. He is currently working toward the master's degree from the Georg August University, Göttingen, Germany.

Marc Timme received the degrees in physics and mathematics from the University of Würzburg, Würzburg, Germany, and the State University of New York at Stony Brook, Stony Brook, NY, USA, and the doctorate degree in theoretical physics from the University of Göttingen, Göttingen, Germany. After two research stays at the Max Planck Institute for Flow Research and the Center for Applied Mathematics, Cornell University, he was appointed to establish a broadly cross-disciplinary research group on Network Dynamics at the Max Planck Institute for Dynamics and Self-Organization. He became an Adjunct Professor with the Institute for Nonlinear Dynamics, the University of Göttingen and is Co-Chair of the Division of Socio-Economic Physics of German Physical Society. He was a Visiting Professor with the TU Darmstadt, Germany, visiting faculty with the ETH Zürich Risk Center and now the Heads the Chair for Network Dynamics as a Strategic Professor with the Institute for Theoretical Physics and the Center for Advancing Electronics Dresden, TU Dresden, Germany. He received a research award of the Berliner Ungewitter Foundation, the Otto Hahn Medal of the Max Planck Society, a Research Fellowship of the National Research Center of Italy and is Honorary Member of Lakeside Labs, Klagenfurt.